

Goat anti-FOXN2 / HTLF Antibody

| | |
|------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Item Number | dAP-0223 |
| Target Molecule | Principle Name: FOXN2 / HTLF; Official Symbol: FOXN2; All Names and Symbols: forkhead box N2; FOXN2; HTLF; human T-cell leukemia virus enhancer factor; T-cell leukemia virus enhancer factor; Accession Number (s): NP_002149.1; Human Gene ID(s): 3344; Non-Human GeneID(s): |
| Immunogen | MCDPLDQLATRTQK, is from N Terminus |
| Applications | Pep ELISA Species Tested: |
| Purification | Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide. |
| Supplied As | lyophilized powder of 50ug or 100ug IgG; Reconstitute IgG with 100ul or 200ul sterile DI Water and final product will be formulated as 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing. |
| Peptide ELISA | Peptide ELISA: antibody detection limit dilution 1 to 16000. |
| Western Blot | Western Blot: No signal obtained yet but low background observed in human lung and Jurkat lysates at up to 1µg/ml. We would appreciate any feedback from people in the field - have any results been reported with other antibodies/lysates? |
| IHC | |
| Reference | Reference(s): Li C, Lusis AJ, Sparkes R, Tran SM, Gaynor R. Characterization and chromosomal mapping of the gene encoding the cellular DNA binding protein HTLF. Genomics. 1992 Jul;13(3):658-64..PMID: 1639393-> |

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for **Research Use Only**