



## Goat anti-Pleckstrin Antibody

<b>Item Number</b>	dAP-0464
<b>Target Molecule</b>	Principle Name: Pleckstrin; Official Symbol: PLEK; All Names and Symbols: PLEK; pleckstrin; P47; p47; FLJ27168; Accession Number (s): NP_002655.2; Human Gene ID(s): 5341; Non-Human GeneID(s):
<b>Immunogen</b>	EWIKAIQMASRTGK, is from C Terminus
<b>Applications</b>	Pep ELISA, WB Species Tested: Human
<b>Purification</b>	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
<b>Supplied As</b>	lyophilized powder of 50ug or 100ug IgG; Reconstitute IgG with 100ul or 200ul sterile DI Water and final product will be formulated as 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
<b>Peptide ELISA</b>	Peptide ELISA: antibody detection limit dilution 1 to 16000.
<b>Western Blot</b>	Western Blot: Approx 45kDa band observed in Human Heart and Peripheral Blood Mononucleocyte lysates (calculated MW of 40.1kDa according to NP_002655.2). Recommended for use at 0.05-1µg/ml.
<b>IHC</b>	
<b>Reference</b>	Reference(s): Tyers M, Rachubinski RA, Stewart MI, Varrichio AM, Shorr RG, Haslam RJ, Harley CB. Molecular cloning and expression of the major protein kinase C substrate of platelets. Nature. 1988 Jun 2;333(6172):470-3..PMID: 2897630 ->

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for **Research Use Only**