

Goat anti-CITED2 (Internal) Antibody

Item Number	dAP-0787
Target Molecule	Principle Name: CITED2 (Internal); Official Symbol: CITED2; All Names and Symbols: CITED2; Cbp/p300-interacting transactivator, with Glu/Asp-rich carboxy-terminal domain, 2 ; HGNC:1987; MRG1; P35SRJ; Accession Number (s): NP_006070.2; Human Gene ID(s): 10370; Non-Human GeneID(s): 17684 (mouse) 114490 (rat)
Immunogen	PNVIDTDFIDEE, is from internal region
Applications	Pep ELISA Species Tested:
Purification	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Supplied As	Lyophilized powder of 50ug or 100ug IgG; Reconstitute IgG with 100ul or 200ul sterile DI Water and final product will be formulated as 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
Peptide ELISA	Peptide ELISA: antibody detection limit dilution 1 to 4000.
Western Blot	Western Blot: Preliminary experiments in adult human heart, kidney, liver and skin lysates gave no specific signal but low background (at antibody concentration up to 1µg/ml). We would appreciate any feedback from people in the field - have any results b
IHC	
Reference	Reference(s): Weninger WJ, Floro KL, Bennett MB, Withington SL, Preis JI, Barbera JP, Mohun TJ, Dunwoodie SL. Cited2 is required both for heart morphogenesis and establishment of the left-right axis in mouse development. Development. 2005 Mar;132(6):1337-48. PMID: 15750185 ->

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the end users! This product is sold for **Research Use Only**