

Goat anti-FPRL1 (aa328-339) Antibody

Item Number	dAP-0828
Target Molecule	Principle Name: FPRL1 (aa328-339); Official Symbol: FPR2; All Names and Symbols: FPR2; FPRL1; ALXR; HM63; FMLPX; FPR2A; FPRH1; FPRH2; LXA4R; FMLP-R-II; formyl peptide receptor-like 1; lipoxin A4 receptor (formyl peptide receptor related); formyl peptide receptor 2; Accession Number (s): NP_001005738.1; NP_001453.1; Human Gene ID(s): 2358; Non-Human GenelD(s):
Immunogen	DSAPTNNTAANS, is from C Terminus Both reported variants (NP_001005738.1 and NP_001453.1) are identical proteins.
Applications	Pep ELISA Species Tested:
Purification	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Supplied As	Lyophilized powder of 50ug or 100ug IgG; Reconstitute IgG with 100ul or 200ul sterile DI Water and final product will be formulated as 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
Peptide ELISA	Peptide ELISA: antibody detection limit dilution 1 to 2000.
Western Blot	Western Blot: Preliminary experiments gave an approx 25kDa band in Human Lymph Node and Tonsil lysates after 1µg/ml antibody staining. Please note that currently we cannot find an explanation in the literature for the band we observe given the calculated
IHC	
Reference	Reference(s): Vachier I, Bonnans C, Chavis C, Farce M, Godard P, Bousquet J, Chanez P. Severe asthma is associated with a loss of LX4, an endogenous anti-inflammatory compound. J Allergy Clin Immunol. 2005 Jan;115(1):55-60.. PMID: 15637547->

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the end users! This product is sold for **Research Use Only**