

## Goat anti-VDAC2 (Internal) Antibody

Item Number	dAP-0898
Target Molecule	Principle Name: VDAC2 (Internal); Official Symbol: VDAC2; All Names and Symbols: VDAC2; voltage-dependent anion channel 2; RP11-375G3.1; FLJ23841 ; Accession Number (s): NP_003366.2; Human Gene ID(s): 7417; Non-Human GenID(s): 22334 (mouse)
Immunogen	DSAKSKLTRNN, is from internal region
Applications	Pep ELISA, WB Species Tested: Human
Purification	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Supplied As	Lyophilized powder of 50ug or 100ug IgG; Reconstitute IgG with 100ul or 200ul sterile DI Water and final product will be formulated as 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
Peptide ELISA	Peptide ELISA: antibody detection limit dilution 1 to 64000.
Western Blot	Western Blot: Approx 32kDa band observed in Human Heart lysates (calculated MW of 31.6 kDa according to NP_003366.2). Recommended concentration: 0.3-1µg/ml.
IHC	
Reference	Reference(s): Chandra D, Choy G, Daniel PT, Tang DG. Bax-dependent regulation of Bak by voltage-dependent anion channel 2. J Biol Chem. 2005 May 13;280(19):19051-61. Epub 2005 Mar 9. PMID: 15757910 ->

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the end users! This product is sold for **Research Use Only**