

Goat anti-VHL Antibody

Item Number	dAP-0917
Target Molecule	Principle Name: VHL; Official Symbol: VHL; All Names and Symbols: VHL; von Hippel-Lindau tumor suppressor; HRCA1; RCA1; VHL1 ; elongin binding protein; von Hippel-Lindau syndrome; Accession Number (s): NP_000542.1; NP_937799.1; Human Gene ID(s): 7428; Non-Human GenelD(s): 22346 (mouse) 24874 (rat)
Immunogen	RSLVKPENYRRRLD, is from internal region This antibody is expected to recognise both reported isoforms (NP_000542.1 and NP_937799.1).
Applications	Pep ELISA, WB, IHC, EIA Species Tested: Human, Mouse, Rat
Purification	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Supplied As	Lyophilized powder of 50ug or 100ug IgG; Reconstitute IgG with 100ul or 200ul sterile DI Water and final product will be formulated as 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
Peptide ELISA	Peptide ELISA: antibody detection limit dilution 1 to 128000.
Western Blot	Western Blot: Approx 17kDa band observed in Human Ovary and Testis lysates and also in lysates of Mouse and Rat Testis and Brain and of all human and mouse cell lines tested (calculated MW of 19.7kDa according to NP_937799.1). Recommended concentration:
IHC	Immunohistochemistry: Paraffin embedded Human Kidney. Recommended concentration: 5µg/ml.
Reference	Reference(s): Kong X, Lin Z, Liang D, Fath D, Sang N, Caro J. Histone deacetylase inhibitors induce VHL and ubiquitin-independent proteasomal degradation of hypoxia-inducible factor 1alpha. Mol Cell Biol. 2006 Mar;26(6):2019-28. PMID: 16507982 ->

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the end users! This product is sold for **Research Use Only**