

## Goat anti-Farnesoid X receptor / FXR Antibody

Item Number	dAP-0963
Target Molecule	Principle Name: Farnesoid X receptor / FXR; Official Symbol: NR1H4; All Names and Symbols: farnesoid X receptor; HRR1; NR1H4; nuclear receptor subfamily 1, group H, member 4 ; BAR; FXR; HRR-1; RIP14 ; farnesol receptor HRR-1; Accession Number (s): NP_005114.1; Human Gene ID(s): 9971; Non-Human GenelD(s):
Immunogen	KSCREKTELTPDQQ, is from internal region
Applications	Pep ELISA, WB Species Tested: Human
Purification	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Supplied As	Lyophilized powder of 50ug or 100ug IgG; Reconstitute IgG with 100ul or 200ul sterile DI Water and final product will be formulated as 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
Peptide ELISA	Peptide ELISA: antibody detection limit dilution 1 to 32000.
Western Blot	Western Blot: Approx 55kDa band observed in Human Duodenum and Human Lung lysates (calculated MW of 54.4kDa according to NP_005114.1). Recommended concentration: 1-3µg/ml.
IHC	
Reference	Reference(s): Huang W, Ma K, Zhang J, Qatanani M, Cuvillier J, Liu J, Dong B, Huang X, Moore DD. Nuclear receptor-dependent bile acid signaling is required for normal liver regeneration. <i>Science</i> . 2006 Apr 14;312(5771):233-6. PMID: 16614213 ->

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the end users! This product is sold for **Research Use Only**