

## Goat anti-Robo3 (mouse) Antibody

<b>Item Number</b>	dAP-1132
<b>Target Molecule</b>	Principle Name: Robo3 (mouse); Official Symbol: Robo3; All Names and Symbols: Robo3; roundabout homolog 3 (Drosophila); Rbig1; RIG1; Rig-1; Robo3a; Robo3b; retinoblastoma inhibiting gene 1; Accession Number (s): NP_035378.1; AAD11628.1; Human Gene ID(s): ; Non-Human GeneID(s): 19649 (mouse) 315564 (rat)
<b>Immunogen</b>	RRQKQRKELSHYT, is from internal region Both variants represent identical protein
<b>Applications</b>	Pep ELISA  Species Tested:
<b>Purification</b>	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
<b>Supplied As</b>	lyophilized powder of 50ug or 100ug IgG; Reconstitute IgG with 100ul or 200ul sterile DI Water and final product will be formulated as 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
<b>Peptide ELISA</b>	Peptide ELISA: antibody detection limit dilution 1 to 16000.
<b>Western Blot</b>	Western Blot: Preliminary experiments gave an approx 75-80kDa band in Mouse Brain and Rat Brain lysates after 0.1µg/ml antibody staining. Please note that currently we cannot find an explanation in the literature for the band we observe given the calcul
<b>IHC</b>	
<b>Reference</b>	Reference(s): Kato H, Takeuchi O, Sato S, Yoneyama M, Yamamoto M, Matsui K, Uematsu S, Jung A, Kawai T, Ishii KJ, Yamaguchi O, Otsu K, Tsujimura T, Koh CS, Reis e Sousa C, Matsuura Y, Fujita T, Akira S. Differential roles of MDA5 and RIG-I helicases in the recognition of RNA viruses. Nature. 2006 May

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for **Research Use Only**