

## Goat anti-MDA5 / Ifih1 (mouse) Antibody

Item Number	dAP-1136
Target Molecule	Principle Name: MDA5 / Ifih1 (mouse); Official Symbol: Ifih1; All Names and Symbols: Ifih1; interferon induced with helicase C domain 1; 9130009C22Rik; Helicard; Hlcd; MDA-5; MDA5; MGC90959; melanoma differentiation associated protein-5; Accession Number (s): NP_082111.1; Human Gene ID(s): ; Non-Human GenID(s): 71586 (mouse)
Immunogen	KQKLRNNDLKKQN, is from internal region
Applications	Pep ELISA Species Tested:
Purification	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Supplied As	Lyophilized powder of 50ug or 100ug IgG; Reconstitute IgG with 100ul or 200ul sterile DI Water and final product will be formulated as 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
Peptide ELISA	Peptide ELISA: antibody detection limit dilution 1 to 16000.
Western Blot	Western Blot: Preliminary experiments gave an approx 80kDa band in Mouse Brain lysates after 1µg/ml antibody staining. Please note that currently we cannot find an explanation in the literature for the band we observe given the calculated size of 116kDa
IHC	
Reference	Reference(s): Kato H, Takeuchi O, Sato S, Yoneyama M, Yamamoto M, Matsui K, Uematsu S, Jung A, Kawai T, Ishii KJ, Yamaguchi O, Otsu K, Tsujimura T, Koh CS, Reis e Sousa C, Matsuura Y, Fujita T, Akira S. Differential roles of MDA5 and RIG-I helicases in the recognition of RNA viruses. <i>Nature</i> . 2006 May

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the end users! This product is sold for **Research Use Only**