

Goat anti-NAT2 (aa182-195) Antibody

Item Number	dAP-1581
Target Molecule	Principle Name: NAT2 (aa182-195); Official Symbol: NAT2 ; All Names and Symbols: NAT2 ; N-acetyltransferase 2 (arylamine N-acetyltransferase) ; AAC2; Arylamine N-acetyltransferase-2; arylamide acetylase 2; arylamide acetylase 2 (N-acetyltransferase 2, isoniazid inactivation); arylamine N-acetyltransferase 2; Accession Number (s): NP_000006.2; Human Gene ID(s): 10; Non-Human GenelD(s):
Immunogen	PKKKHQKIQYLFTLE, is from internal region
Applications	Pep ELISA Species Tested:
Purification	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Supplied As	Lyophilized powder of 50ug or 100ug IgG; Reconstitute IgG with 100ul or 200ul sterile DI Water and final product will be formulated as 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
Peptide ELISA	Peptide ELISA: antibody detection limit dilution 1 to 16000.
Western Blot	Western Blot: Preliminary experiments gave an approx 27 kDa band in Human Kidney, Liver and Testis lysates after 1µg/ml antibody staining. Please note that currently we cannot find an explanation in the literature for the band we observe given the calcul
IHC	
Reference	Reference(s): Wu H, Dombrovsky L, Tempel W, Martin F, Loppnau P, Goodfellow GH, Grant DM, Plotnikov AN. Structural basis of substrate-binding specificity of human arylamine N-acetyltransferases. <i>J Biol Chem.</i> 2007 Oct 12;282(41):30189-97. Epub 2007 Jul 26..PMID: 17656365 ->

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the end users! This product is sold for **Research Use Only**