



## Goat anti-Ankyrin 1 / ANK1 Antibody

<b>Item Number</b>	dAP-1591
<b>Target Molecule</b>	Principle Name: Ankyrin 1 / ANK1; Official Symbol: ANK1 ; All Names and Symbols: ANK1 ; ankyrin 1, erythrocytic ; ANK; SPH1; SPH2; ankyrin 1; ankyrin-1, erythrocytic; ankyrin-R; Accession Number (s): NP_000028.3; NP_065208.2; NP_065209.2; NP_065210.2; NP_065211.2; NP_065213.2; Human Gene ID (s): 286; Non-Human GeneID(s): 11733 (mouse) 306570 (rat)
<b>Immunogen</b>	QIVKRASLKRKQ, is from C Terminus This antibody is expected to recognise isoform 3 (NP_000028.3) and isoform 5 (NP_065211.2).
<b>Applications</b>	Pep ELISA  Species Tested:
<b>Purification</b>	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
<b>Supplied As</b>	lyophilized powder of 50ug or 100ug IgG; Reconstitute IgG with 100ul or 200ul sterile DI Water and final product will be formulated as 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
<b>Peptide ELISA</b>	Peptide ELISA: antibody detection limit dilution 1 to 128000.
<b>Western Blot</b>	Western Blot: Preliminary experiments gave approx. 19-23 kDa bands in Human Skeletal Muscle lysates after 0.01µg/ml antibody staining. Please note that currently we cannot find an explanation in the literature for these bands we observe given the observe
<b>IHC</b>	
<b>Reference</b>	Reference(s): Dubreuil RR. Functional links between membrane transport and the spectrin cytoskeleton. J Membr Biol. 2006;211(3):151-61. Epub 2006 Nov 7. Review..PMID: 17091212 ->

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for **Research Use Only**