



Goat anti-CCL3L1 Antibody

Item Number	dAP-1628
Target Molecule	Principle Name: CCL3L1; Official Symbol: CCL3L1; All Names and Symbols: CCL3L1; chemokine (C-C motif) ligand 3-like 1; 464.2; D17S1718; G0S19-2; LD78; LD78BETA; MGC104178; MGC12815; SCYA3L1; SCYA3L1; small inducible cytokine A3-like 1; Accession Number (s): NP_066286.1; Human Gene ID(s): 6349; Non-Human GeneID(s):
Immunogen	CNQVLSAPLAADT , is from internal region
Applications	Pep ELISA Species Tested:
Purification	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Supplied As	lyophilized powder of 50ug or 100ug IgG; Reconstitute IgG with 100ul or 200ul sterile DI Water and final product will be formulated as 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
Peptide ELISA	Peptide ELISA: antibody detection limit dilution 1 to 8000.
Western Blot	Western Blot: Preliminary experiments in Human Bone Marrow and in Peripheral Blood Mononucleocyte lysates gave no specific signal but low background (at antibody concentration up to 1µg/ml). We would appreciate any feedback from people in the field - hav
IHC	
Reference	Reference(s): McKinney C, Merriman ME, Chapman PT, Gow PJ, Harrison AA, Highton J, Jones PB, McLean L, O'Donnell JL, Pokorny V, Spellerberg M, Stamp LK, Willis J, Steer S, Merriman TR. Evidence for an influence of chemokine ligand 3-like 1 (CCL3L1) gene copy number on sus-

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for **Research Use Only**