

## Goat anti-ZNF228 / ZFP112 Antibody

<b>Item Number</b>	dAP-1673
<b>Target Molecule</b>	Principle Name: ZNF228 / ZFP112; Official Symbol: ZFP112; All Names and Symbols: ZFP112; zinc finger protein 112 homolog (mouse); ZNF228; zinc finger protein 228; Accession Number (s): NP_001076804.1; NP_037512.3; Human Gene ID(s): 7771; Non-Human GeneID(s):
<b>Immunogen</b>	EREDDIENSHLKSQY , is from internal region This antibody is expected to recognize both reported isoforms (NP_001076804.1; NP_037512.3)
<b>Applications</b>	Pep ELISA  Species Tested:
<b>Purification</b>	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
<b>Supplied As</b>	lyophilized powder of 50ug or 100ug IgG; Reconstitute IgG with 100ul or 200ul sterile DI Water and final product will be formulated as 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
<b>Peptide ELISA</b>	Peptide ELISA: antibody detection limit dilution 1 to 4000.
<b>Western Blot</b>	Western Blot: Preliminary experiments gave an approx. 48kDa band in lysates of cell line HepG2 after 2µg/ml antibody staining. Please note that currently we cannot find an explanation in the literature for the band we observe given the calculated size of
<b>IHC</b>	
<b>Reference</b>	Reference(s): Shannon, M., Hamilton, A.T., Gordon, L., Branscomb, E. and Stubbs, L. Differential expansion of zinc-finger transcription factor loci in homologous human and mouse gene clusters. Genome Res. 13 (6A), 1097-1110 (2003)..PMID: 12743021->

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for **Research Use Only**