

## Goat anti-CAGE1 Antibody

<b>Item Number</b>	dAP-1787
<b>Target Molecule</b>	Principle Name: CAGE1; Official Symbol: CAGE1; All Names and Symbols: CAGE1; cancer antigen 1; CTAG3; FLJ40441; bA69L16.7; cancer-associated gene 1; cancer/testis antigen 3; cancer/testis antigen gene 1; Accession Number (s): NP_995586.1; Human Gene ID(s): 285782; Non-Human GeneID(s):
<b>Immunogen</b>	ARKPRSKSLENHPKS, is from internal region
<b>Applications</b>	Pep ELISA Species Tested:
<b>Purification</b>	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
<b>Supplied As</b>	lyophilized powder of 50ug or 100ug IgG; Reconstitute IgG with 100ul or 200ul sterile DI Water and final product will be formulated as 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
<b>Peptide ELISA</b>	Peptide ELISA: antibody detection limit dilution 1 to 16000.
<b>Western Blot</b>	Western Blot: Preliminary experiments gave an approx. 37kDa band in Human Ovary, Testis and Thyroid Gland lysates after 0.5µg/ml antibody staining. Please note that currently we cannot find an explanation in the literature for the band we observe given t
<b>IHC</b>	
<b>Reference</b>	Reference(s): Kunze E, Schlott T. High frequency of promoter methylation of the 14-3-3 sigma and CAGE-1 genes, but lack of hypermethylation of the caveolin-1 gene, in primary adenocarcinomas and signet ring cell carcinomas of the urinary bladder. International journal of molecular medicine 2007 Oct 20 (4): 557-

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for **Research Use Only**