

## Goat anti-UPF3A / RENT3A Antibody

<b>Item Number</b>	dAP-1933
<b>Target Molecule</b>	Principle Name: UPF3A / RENT3A; Official Symbol: UPF3A; All Names and Symbols: UPF3A; UPF3 regulator of nonsense transcripts homolog A (yeast); HUPF3A; RENT3A; UPF3; OTTHUMP00000018789; UPF3 regulator of nonsense transcripts homolog A; Accession Number (s): NP_075387.1; NP_542418.1; Human Gene ID(s): 65110; Non-Human GenelD(s):
<b>Immunogen</b>	KETDKQKKIAEKE, is from internal region This antibody is expected to recognize both isoforms (NP_075387.1; NP_542418.1).
<b>Applications</b>	Pep ELISA  Species Tested:
<b>Purification</b>	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
<b>Supplied As</b>	Lyophilized powder of 50ug or 100ug IgG; Reconstitute IgG with 100ul or 200ul sterile DI Water and final product will be formulated as 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
<b>Peptide ELISA</b>	Peptide ELISA: antibody detection limit dilution 1 to 8000.
<b>Western Blot</b>	Western Blot: Preliminary experiments gave an approx 48kDa band in lysates of cell lines HeLa and Kelly after 0.3µg/ml antibody staining. Please note that currently we cannot find an explanation in the literature for the band we observe given the calcula
<b>IHC</b>	
<b>Reference</b>	Reference(s): Tarpey PS, Raymond FL, Nguyen LS, Rodriguez J, Hackett A, Vandeleur L, Smith R, Shoubridge C, Edkins S, Stevens C, O'Meara S, Tofts C, Bartherope S, Buck G, Cole J, Halliday K, Hills K, Jones D, Mironenko T, Perry J, Varian J, West S, Widaa S, Teague J, Di Mutations in UPF3B, a member of

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the end users! This product is sold for **Research Use Only**