

## Goat anti-HIP1R/ HIP12 Antibody

<b>Item Number</b>	dAP-1989
<b>Target Molecule</b>	Principle Name: HIP1R/ HIP12; Official Symbol: HIP1R; All Names and Symbols: HIP1R; huntingtin interacting protein 1 related; FLJ14000; HIP12; HIP3; ILWEQ; KIAA0655; MGC47513; huntingtin interacting protein 12; huntingtin interacting protein-1-related; Accession Number (s): NP_003950.1; Human Gene ID (s): 9026; Non-Human GeneID(s):
<b>Immunogen</b>	RQDHQLDKKDG IYP, is from C Terminus
<b>Applications</b>	Pep ELISA  Species Tested:
<b>Purification</b>	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
<b>Supplied As</b>	lyophilized powder of 50ug or 100ug IgG; Reconstitute IgG with 100ul or 200ul sterile DI Water and final product will be formulated as 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
<b>Peptide ELISA</b>	Peptide ELISA: antibody detection limit dilution 1 to 32000.
<b>Western Blot</b>	Western Blot: Preliminary experiments in Human Brain (Amygdala, Hippocampus and Substantia nigra) lysates gave no specific signal but low background (at antibody concentration up to 1µg/ml). We would appreciate any feedback from people in the field - hav
<b>IHC</b>	

<b>Reference</b>	Reference(s): Poupon V, Girard M, Legendre-Guillemin V, Thomas S, Bourbonniere L, Philie J, Bright NA, McPherson PS, Clathrin light chains function in mannose phosphate receptor trafficking via regulation of actin assembly. Proc. Natl. Acad. Sci. U.S.A. 2008 Jan 105 (1): 168-73..PMID: 18165318->
------------------	--

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for **Research Use Only**