

Goat anti-HOXD13 Antibody

Item Number	dAP-2064
Target Molecule	Principle Name: HOXD13; Official Symbol: HOXD13; All Names and Symbols: HOXD13; homeobox D13; BDE; BDSD; HOX4I; SPD; homeo box 4I; homeo box D13; Accession Number (s): NP_000514.2; Human Gene ID(s): 3239; Non-Human GenelD(s): 15433 (mouse) 288154 (rat)
Immunogen	KSSFPGDVALNQPD, is from internal region
Applications	Pep ELISA, IHC Species Tested: Human
Purification	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Supplied As	Lyophilized powder of 50ug or 100ug IgG; Reconstitute IgG with 100ul or 200ul sterile DI Water and final product will be formulated as 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
Peptide ELISA	Peptide ELISA: antibody detection limit dilution 1 to 8000.
Western Blot	Western Blot: Preliminary experiments in Human Breast cancer lysates gave no specific signal but low background (at antibody concentration up to 1µg/ml). We would appreciate any feedback from people in the field - have any results been reported with othe
IHC	Immunohistochemistry: In paraffin embedded Human Prostate shows nuclear staining in the secretory cells. Recommended concentration, 1-2µg/ml.
Reference	Reference(s): Cantile M, Franco R, Tschan A, Baumhoer D, Zlobec I, Schiavo G, Forte I, Bihl M, Ligueri G, Botti G, Tornillo L, Karamitopoulou-Diamantis E, Terracciano L, Cillo C, HOX D13 expression across 79 tumor tissue types. Int J Cancer. 2009 Oct 1;125(7):1532-41..PMID: 19488988-->

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for **Research Use Only**