

## Goat anti-CDC2L6 Antibody

<b>Item Number</b>	dAP-2256
<b>Target Molecule</b>	Principle Name: CDC2L6; Official Symbol: CDC2L6; All Names and Symbols: CDC2L6; cell division cycle 2-like 6 (CDK8-like); CDK11; KIAA1028; bA346C16.3; CDK8-like cyclin-dependent kinase; cyclin-dependent kinase (CDC2-like) 11; death-preventing kinase; Accession Number (s): NP_055891.1; Human Gene ID(s): 23097; Non-Human GeneID(s): 78334 (mouse) 309804 (rat)
<b>Immunogen</b>	QYHPSHQAH, is from C Terminus
<b>Applications</b>	Pep ELISA Species Tested:
<b>Purification</b>	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
<b>Supplied As</b>	Lyophilized powder of 50ug or 100ug IgG; Reconstitute IgG with 100ul or 200ul sterile DI Water and final product will be formulated as 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
<b>Peptide ELISA</b>	Peptide ELISA: antibody detection limit dilution 1 to 8000.
<b>Western Blot</b>	Western Blot: Preliminary experiments gave bands at approx 45kDa and 20kDa in Rat Eye lysates after 1µg/ml antibody staining. Please note that currently we cannot find an explanation in the literature for the bands we observe given the calculated size of
<b>IHC</b>	
<b>Reference</b>	Reference(s): Malumbres M, Harlow E, Hunt T, Hunter T, Lahti JM, Manning G, Morgan DO, Tsai LH, Wolgemuth DJ. Cyclin-dependent kinases: a family portrait. <i>Nat Cell Biol.</i> 2009 Nov;11(11):1275-6..PMID: 19884882->

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the end users! This product is sold for **Research Use Only**