

Goat anti-SDCCAG8 (aa469-483) Antibody

Item Number	dAP-2629
Target Molecule	Principle Name: SDCCAG8 (aa469-483); Official Symbol: SDCCAG8; All Names and Symbols: SDCCAG8; serologically defined colon cancer antigen 8; BBS16; CCCAP; NPHP10; NY-CO-8; SLSN7; hCCCAP; OT-THUMP00000037913; antigen NY-CO-8; centrosomal colon cancer autoantigen protein; Accession Number (s): NP_006633.1; Human Gene ID(s): 10806; Non-Human GeneID(s): 76816 (mouse) 305002 (rat)
Immunogen	DEAEKEHREFRAKTN, is from internal region
Applications	Pep ELISA Species Tested:
Purification	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Supplied As	lyophilized powder of 50ug or 100ug IgG; Reconstitute IgG with 100ul or 200ul sterile DI Water and final product will be formulated as 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
Peptide ELISA	Peptide ELISA: antibody detection limit dilution 1 to 32000.
Western Blot	Western Blot: Preliminary experiments in Human Prostate, Tonsil and Thymus lysates gave no specific signal but low background (at antibody concentration up to 1µg/ml). We would appreciate any feedback from people in the field - have any results been repo
IHC	
Reference	Reference(s): Otto EA, Hurd TW, Airik R, Chaki M, Zhou W, Stoetzel C, Patil SB, Levy S, Ghosh AK, Murga-Zamalloa CA, van Reeuwijk J, Letteboer SJ, Sang L, Giles RH, Liu Q, Coene KL, Estrada-Cuzcano A, Collin RW, McLaughlin HM, Held S, Kasanuki JM, Ramaswami G, Conte J Candidate exome capture iden-

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for **Research Use Only**