

## Goat anti-STR8 (aa70-81) Antibody

Item Number	dAP-2654
Target Molecule	Principle Name: STRA8 (aa70-81); Official Symbol: STRA8; All Names and Symbols: STRA8; stimulated by retinoic acid gene 8 homolog (mouse); OTTHUHMP00000207988; stimulated by retinoic acid gene 8 protein homolog; Accession Number (s): NP_872295.1; Human Gene ID(s): 346673; Non-Human GenelD(s): 500079 (rat)
Immunogen	NNLRKTVYSQSD, is from internal region
Applications	Pep ELISA Species Tested:
Purification	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Supplied As	Lyophilized powder of 50ug or 100ug IgG; Reconstitute IgG with 100ul or 200ul sterile DI Water and final product will be formulated as 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
Peptide ELISA	Peptide ELISA: antibody detection limit dilution 1 to 8000.
Western Blot	Western Blot: Preliminary experiments gave an approx 90kDa band in Rat Duodenum and Testis lysates after 1µg/ml antibody staining. Please note that currently we cannot find an explanation in the literature for the band we observe given the calculated size.
IHC	
Reference	Reference(s): Le Bouffant R, Guerquin MJ, Duquenne C, Frydman N, Coffigny H, Rouiller Fabre V, Frydman R, Habert R, Livera G. Meiosis initiation in the human ovary requires intrinsic retinoic acid synthesis. Hum Reprod. 2010 Oct;25(10):2579-90.. PMID: 20670969->

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the end users! This product is sold for **Research Use Only**