

Goat anti-PCNA (aa111-122) Antibody

Item Number	dAP-2699
Target Molecule	Principle Name: PCNA (aa111-122); Official Symbol: PCNA; All Names and Symbols: PCNA; proliferating cell nuclear antigen; MGC8367; DNA polymerase delta auxiliary protein; OTTHUMP00000030189; OTTHUMP00000030190; cyclin; Accession Number (s): NP_002583.1; Human Gene ID(s): 5111; Non-Human GeneID(s): 18538 (mouse) 25737 (rat)
Immunogen	EAPNQEKVSDYE, is from internal region Reported variants represent identical protein: NP_872590.1, NP_002583.1
Applications	Pep ELISA, WB, EIA Species Tested: Human, Mouse, Rat, Pig
Purification	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Supplied As	lyophilized powder of 50ug or 100ug IgG; Reconstitute IgG with 100ul or 200ul sterile DI Water and final product will be formulated as 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
Peptide ELISA	Peptide ELISA: antibody detection limit dilution 1 to 128000.
Western Blot	Western Blot: Approx 30kDa band observed in lysates of cell line Jurkat (calculated MW of 28.8kDa according to Human NP_002583.1). Approx.35kDa band was observed in lysates of cell line NIH3T3 and in Mouse and Rat Testis, and approx 33kDa in Pig Spleen
IHC	
Reference	Reference(s): Jones MJ, Colnaghi L, Huang TT. Dysregulation of DNA polymerase κ recruitment to replication forks results in genomic instability. EMBO J. 2011 Dec 13..PMID: 22157819->

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for **Research Use Only**