



## Goat anti-NEUROD2 (aa23-37) Antibody

<b>Item Number</b>	dAP-2788
<b>Target Molecule</b>	Principle Name: NEUROD2 (aa23-37); Official Symbol: NEUROD2; All Names and Symbols: NEUROD2; neurogenic differentiation 2; MGC26304; NDRF; bHLHa1; class A basic helix-loop-helix protein 1; neuroD-related factor; neurogenic basic-helix-loop-helix protein; neurogenic differentiation factor 2; Accession Number (s): NP_006151.3; Human Gene ID(s): 4761; Non-Human GeneID(s):
<b>Immunogen</b>	DGEDDEPRSDKGDAP, is from internal region (near N Terminus)
<b>Applications</b>	Pep ELISA Species Tested:
<b>Purification</b>	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
<b>Supplied As</b>	lyophilized powder of 50ug or 100ug IgG; Reconstitute IgG with 100ul or 200ul sterile DI Water and final product will be formulated as 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
<b>Peptide ELISA</b>	Peptide ELISA: antibody detection limit dilution 1 to 2000.
<b>Western Blot</b>	Western Blot: Preliminary experiments in Human Brain lysates from different regions gave no specific signal but low background (at antibody concentration up to 1µg/ml). We would appreciate any feedback from people in the field - have any results been rep
<b>IHC</b>	
<b>Reference</b>	Reference(s): Zill P, Büttner A, Eisenmenger W, Bondy B. A possible impact of the neuroD2 transcription factor on the development of drug abusing behavior. Mol Psychiatry. 2006 Jun;11(6):525-7..PMID: 16607387->

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for **Research Use Only**