

Goat anti-MAPKAPK2 (aa44-55) Antibody

Item Number	dAP-2870
Target Molecule	Principle Name: MAPKAPK2 (aa44-55); Official Symbol: MAPKAPK2; All Names and Symbols: MAPKAP-K2; mitogen-activated protein kinase-activated protein kinase 2; MAPKAP-K2; MK-2; MK2; MAP kinase-activated protein kinase 2; MAPK-activated protein kinase 2; MAPKAP kinase 2; Accession Number (s): NP_004750.1; NP_116584.2; Human Gene ID(s): 9261; Non-Human GeneID(s): 17164 (mouse) 289014 (rat)
Immunogen	PQFHVKSGLQIK, is from internal region This antibody is expected to recognize both reported isoforms (NP_004750.1; NP_116584.2).
Applications	Pep ELISA, WB Trf, IP Species Tested: Moiuse
Purification	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Supplied As	lyophilized powder of 50ug or 100ug IgG; Reconstitute IgG with 100ul or 200ul sterile DI Water and final product will be formulated as 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
Peptide ELISA	Peptide ELISA: antibody detection limit dilution 1 to 32000.
Western Blot	Western Blot: In transfected HEK293 transiently expressing either Mouse MK2 fused to GFP or the similar Mouse MK5 fused to GFP, only the expected band was observed in the MK2 lane after EB11929 and EB11930 labeling. Data obtained from Dr. M. B. Menon, I
IHC	
Reference	Reference(s): Yang Y, Xia F, Hermance N, Mabb A, Simonson S, Morrissey S, Gandhi P, Munson M, Miyamoto S, Kelliher MA. A cytosolic ATM/NEMO/RIP1 complex recruits TAK1 to mediate the NF- κ B and p38 mitogen-activated protein kinase (MAPK)/MAPK-activated protein 2 responses to DNA damage. Mol

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for **Research Use Only**