

Goat anti-Osteocalcin Antibody

| | |
|-----------------|---|
| Item Number | dAP-2881 |
| Target Molecule | Principle Name: Osteocalcin; Official Symbol: BGLAP; All Names and Symbols: BGLAP; bone gamma-carboxyglutamate (gla) protein; BGP; OC; OTTHUMP0000016586; bone Gla protein; bone gamma-carboxyglutamate (gla) protein (osteocalcin); gamma-carboxyglutamic acid-containing protein; osteocalcin; Accession Number (s): NP_954642.1; Human Gene ID(s): 632; Non-Human GeneID(s): |
| Immunogen | DELADHIGFQEAYRR, is from internal region (near C terminus) |
| Applications | Pep ELISA Species Tested: |
| Purification | Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide. |
| Supplied As | Lyophilized powder of 50ug or 100ug IgG; Reconstitute IgG with 100ul or 200ul sterile DI Water and final product will be formulated as 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing. |
| Peptide ELISA | Peptide ELISA: antibody detection limit dilution 1 to 128000. |
| Western Blot | Western Blot: Not yet tested - our routinely used western blotting protocol does not allow detection of proteins smaller than the calculated size of 11.0kDa according to the precursor NP_954642.1. Therefore we cannot recommend an optimal concentration an |
| IHC | |
| Reference | Reference(s): Iglesias P, Arrieta F, Piñera M, Botella-Carretero JI, Balsa JA, Zamarrón I, Menacho M, Díez JJ, Muñoz T, Vázquez C. Serum concentrations of osteocalcin, procollagen type 1 N-terminal propeptide and beta-CrossLaps in obese subjects with varying degrees of glucose tolerance. Clin Endocrinol (Oxf). |

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the end users! This product is sold for **Research Use Only**