

Goat anti-FURIN / PCSK3 Antibody

Item Number	dAP-2898
Target Molecule	Principle Name: FURIN / PCSK3; Official Symbol: FURIN; All Names and Symbols: FURIN; furin (paired basic amino acid cleaving enzyme); FUR; PACE; PCSK3; SPC1; FES upstream region; dibasic processing enzyme; furin; furin, membrane associated receptor protein; paired basic amino acid cleaving enzyme (furin, membrane associated receptor); Accession Number (s): NP_002560.1; Human Gene ID(s): 5045; Non-Human GenID(s):
Immunogen	RTKRDVYQEPTDP, is from internal region
Applications	Pep ELISA, WB Trf Species Tested: Human
Purification	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Supplied As	Lyophilized powder of 50ug or 100ug IgG; Reconstitute IgG with 100ul or 200ul sterile DI Water and final product will be formulated as 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
Peptide ELISA	Peptide ELISA: antibody detection limit dilution 1 to 8000.
Western Blot	Western Blot: In transfected HEK293 transiently expressing Human Furin (myc and DYKDDDDK tagged), a band of approx. 110kDa is observed. No bands are observed in mock-transfected HEK293 and the same band is observed using anti-myc antibody. Recommended c
IHC	
Reference	Reference(s): Jin W, Wang X, Millar JS, Quertermous T, Rothblat GH, Glick JM, Rader DJ Hepatic proprotein convertases modulate HDL metabolism Cell Metab. 2007 Aug;6(2):129-36. PMID: 17681148->

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the end users! This product is sold for **Research Use Only**