

Goat anti-IFNAR1 Antibody

Item Number	dAP-2913
Target Molecule	Principle Name: IFNAR1; Official Symbol: IFNAR1; All Names and Symbols: IFNAR1; interferon (alpha, beta and omega) receptor 1; AVP; IFN-alpha-REC; IFNAR; IFNBR; IFRC; CRF2-1; IFN-R-1; IFN-alpha/beta receptor 1; alpha-type antiviral protein; beta-type antiviral protein; cytokine receptor class-II member 1; cytokine receptor fa; Accession Number (s): NP_000620.2; Human Gene ID(s): 3454; Non-Human GenelD (s):
Immunogen	KQIPDCENVKTTQ, is from internal region
Applications	Pep ELISA Species Tested:
Purification	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Supplied As	Lyophilized powder of 50ug or 100ug IgG; Reconstitute IgG with 100ul or 200ul sterile DI Water and final product will be formulated as 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
Peptide ELISA	Peptide ELISA: antibody detection limit dilution 1 to 128000.
Western Blot	Western Blot: Preliminary experiments in lysates of cell lines A431, K562 and U937 gave no specific signal but low background (at antibody concentration up to 1µg/ml). We would appreciate any feedback from people in the field - have any results been reported?
IHC	
Reference	Reference(s): Kurche JS, Haluszczak C, McWilliams JA, Sanchez PJ, Kedl RM. Type I IFN-dependent T cell activation is mediated by IFN-dependent dendritic cell OX40 ligand expression and is independent of T cell IFNAR expression. <i>J Immunol.</i> 2012 Jan 15;188(2):585-93.. PMID: 22156349->

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the end users! This product is sold for **Research Use Only**