

Goat anti-BATF Antibody

Item Number	dAP-2921
Target Molecule	Principle Name: BATF; Official Symbol: BATF; All Names and Symbols: BATF; basic leucine zipper transcription factor, ATF-like; B-ATF; BATF1; SFA-2; SFA2; B-cell-activating transcription factor; SF-HT-activated gene 2 protein; activating transcription factor B; basic leucine zipper transcriptional factor ATF-like; Accession Number (s): NP_006390.1; Human Gene ID(s): 10538; Non-Human GenelD(s): 53314 (mouse) 299206 (rat)
Immunogen	EVVYSAHAFHQPHVS, is from C Terminus
Applications	Pep ELISA Species Tested:
Purification	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Supplied As	lyophilized powder of 50ug or 100ug IgG; Reconstitute IgG with 100ul or 200ul sterile DI Water and final product will be formulated as 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
Peptide ELISA	Peptide ELISA: antibody detection limit dilution 1 to 128000.
Western Blot	Western Blot: Preliminary experiments gave an approx 25kDa band in Human Spleen lysates after 1µg/ml antibody staining. Please note that currently we cannot find an explanation in the literature for the band we observe given the calculated size of 14.1kD
IHC	
Reference	Reference(s): Quigley M, Pereyra F, Nilsson B, Porichis F, Fonseca C, Eichbaum Q, Julg B, Jesneck JL, Brosnahan K, Imam S, Russell K, Toth I, Piechocka-Trocha A, Dolfi D, Angelosanto J, Crawford A, Shin H, Kwon DS, Zupkosky J, Francisco L, Freeman GJ, Wherry EJ, Kaufma Transcriptional analysis of HIV-

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for **Research Use Only**