

## Goat anti-FBXO4 Antibody

<b>Item Number</b>	dAP-3006
<b>Target Molecule</b>	Principle Name: FBXO4; Official Symbol: FBXO4; All Names and Symbols: FBXO4; F-box protein 4; DKFZp547N213; FBX4; FLJ10141; F-box only protein 4; F-box protein Fbx4; Accession Number (s): NP_036308.1; NP_277019.1; Human Gene ID(s): 26272; Non-Human GenelD(s): 106052 (mouse) 310363 (rat)
<b>Immunogen</b>	PTAGLPQRQIDG, is from internal region This antibody is expected to recognize isoform 1 and 2 (NP_036308.1; NP_277019.1).
<b>Applications</b>	Pep ELISA, WB  Species Tested: Human
<b>Purification</b>	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
<b>Supplied As</b>	Lyophilized powder of 50ug or 100ug IgG; Reconstitute IgG with 100ul or 200ul sterile DI Water and final product will be formulated as 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
<b>Peptide ELISA</b>	Peptide ELISA: antibody detection limit dilution 1 to 16000.
<b>Western Blot</b>	Western Blot: Approx 35kDa band observed in Human Testis lysates (calculated MW of 35kDa according to NP_277019.1). Recommended concentration: 0.5-1.5µg/ml.
<b>IHC</b>	
<b>Reference</b>	Reference(s): Barbash O, Zamfirova P, Lin DI, Chen X, Yang K, Nakagawa H, Lu F, Rustgi AK, Diehl JA, Mutations in Fbx4 inhibit dimerization of the SCF(Fbx4) ligase and contribute to cyclin D1 overexpression in human cancer. <i>Cancer cell</i> 2008 Jul 14 (1): 68-78.. PMID: 18598945->

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for **Research Use Only**