

## Goat anti-JAG1 Antibody

|                        |  |
|------------------------|--|
| <b>Item Number</b>     | dAP-3014   |
| <b>Target Molecule</b> | Principle Name: JAG1; Official Symbol: JAG1; All Names and Symbols: JAG1; jagged 1; AGS; AHD; AWS; CD339; HJ1; JAGL1; MGC104644; protein jagged-1; Accession Number (s): NP_000205.1; Human Gene ID(s): 182; Non-Human GeneID(s): 16449 (mouse) 29146 (rat)                              |
| <b>Immunogen</b>       | TNKQDNRDLESAQS, is from C Terminus   |
| <b>Applications</b>    | Pep ELISA, IHC<br>Species Tested: Human  |
| <b>Purification</b>    | Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.  |
| <b>Supplied As</b>     | Lyophilized powder of 50ug or 100ug IgG; Reconstitute IgG with 100ul or 200ul sterile DI Water and final product will be formulated as 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.<br>Aliquot and store at -20°C. Minimize freezing and thawing. |
| <b>Peptide ELISA</b>   | Peptide ELISA: antibody detection limit dilution 1 to 8000.  |
| <b>Western Blot</b>    |  |
| <b>IHC</b>             | Immunohistochemistry: In paraffin embedded Human Kidney shows membrane staining in the epithelial cells of convoluted tubules. Recommended concentration, 2-4µg/ml.  |
| <b>Reference</b>       | Reference(s): Shimizu M, Cohen B, Goldvasser P, Berman H, Virtanen C, Reedijk M. Plasminogen activator uPA is a direct transcriptional target of the JAG1-Notch receptor signaling pathway in breast cancer. <i>Cancer Res.</i> 2011 Jan 1;71(1):277-86..PMID: 21199807->                |

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the end users! This product is sold for **Research Use Only**