

## Goat anti-Contactin 6 / NB-3 (mouse aa160-172) Antibody

<b>Item Number</b>	dAP-3148
<b>Target Molecule</b>	Principle Name: Contactin 6 / NB-3 (mouse aa160-172); Official Symbol: Cntn6; All Names and Symbols: Cntn6; contactin 6; NB-3; contactin-6; mNB-3; neural recognition molecule NB-3; Accession Number (s): NP_059079.2; Human Gene ID(s): 27255; Non-Human GeneID(s): 53870 (mouse)
<b>Immunogen</b>	NDSPLYVQEDKRR, is from internal region
<b>Applications</b>	Pep ELISA Species Tested:
<b>Purification</b>	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
<b>Supplied As</b>	lyophilized powder of 50ug or 100ug IgG; Reconstitute IgG with 100ul or 200ul sterile DI Water and final product will be formulated as 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
<b>Peptide ELISA</b>	Peptide ELISA: antibody detection limit dilution 1 to 128000.
<b>Western Blot</b>	Western Blot: Preliminary experiments gave an approx 200kDa band in Mouse fetal Brain lysates after 1µg/ml antibody staining. Please note that currently we cannot find an explanation in the literature for the band we observe given the calculated size of
<b>IHC</b>	
<b>Reference</b>	Reference(s): Sakurai K, Toyoshima M, Takeda Y, Shimoda Y, Watanabe K. Synaptic formation in subsets of glutamatergic terminals in the mouse hippocampal formation is affected by a deficiency in the neural cell recognition molecule NB-3. Neurosci Lett. 2010 Apr 5;473(2):102-6..PMID: 20176085->

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for **Research Use Only**