



## Goat anti-NR5A2 / LRH1 (aa328-338) Antibody

<b>Item Number</b>	dAP-3301
<b>Target Molecule</b>	Principle Name: NR5A2 / LRH1 (aa328-338); Official Symbol: NR5A2; All Names and Symbols: NR5A2; nuclear receptor subfamily 5, group A, member 2; B1F; B1F2; CPF; FTF; FTZ-F1; FTZ-F1beta; LRH-1; LRH1; hB1F-2; CYP7A promoter-binding factor; alpha-1-fetoprotein transcription factor; b1-binding factor, hepatocyte transcription factor which activates; Accession Number (s): NP_995582.1; NP_003813.1; Human Gene ID(s): 2494; Non-Human GeneID(s):
<b>Immunogen</b>	QEQANRSKHEK, is from internal region This antibody is expected to recognize both reported isoforms (NP_995582.1; NP_003813.1).
<b>Applications</b>	Pep ELISA, WB  Species Tested: Human
<b>Purification</b>	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
<b>Supplied As</b>	lyophilized powder of 50ug or 100ug IgG; Reconstitute IgG with 100ul or 200ul sterile DI Water and final product will be formulated as 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
<b>Peptide ELISA</b>	Peptide ELISA: antibody detection limit dilution 1 to 128000.
<b>Western Blot</b>	Western Blot: Approx 55kDa band observed in lysates of cell line HepG2 (calculated MW of 56.5kDa according to NP_003813.1). Recommended concentration: 1-3µg/ml.
<b>IHC</b>	
<b>Reference</b>	Reference(s): Yumoto F, Nguyen P, Sablin EP, Baxter JD, Webb P, Fletterick RJ. Structural basis of coactivation of liver receptor homolog-1 by $\beta$ -catenin. Proc Natl Acad Sci U S A. 2012 Jan 3;109(1):143-8..PMID: 22187462->

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for **Research Use Only**