

## Goat anti-Latexin, Biotinylated Antibody

<b>Item Number</b>	dAP-3344
<b>Target Molecule</b>	Principle Name: Latexin, Biotinylated; Official Symbol: LXN; All Names and Symbols: LXN; latexin; ECI; TCI; MUM; endogenous carboxypeptidase inhibitor; tissue carboxypeptidase inhibitor; Accession Number (s): NP_064554.3; Human Gene ID(s): 56925; Non-Human GeneID(s): 17035 (mouse) 59073 (rat)
<b>Immunogen</b>	QYGTKVKHNSRLPKE., is from C Terminus
<b>Applications</b>	Pep ELISA, WB, IHC, IF Species Tested: Human, Mouse, Rat, Dog
<b>Purification</b>	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
<b>Supplied As</b>	lyophilized powder of 50ug or 100ug IgG; Reconstitute IgG with 100ul or 200ul sterile DI Water and final product will be formulated as 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
<b>Peptide ELISA</b>	Peptide ELISA: antibody detection limit dilution 1 to 128000.
<b>Western Blot</b>	Western Blot: Approx 28kDa band observed in Rat Brain lysates (calculated MW of 25.6kDa according to Rat NP_113843.1). See non-biotinylated parental product's datasheet for further QC data. Recommended concentration: 0.1-0.3µg/ml.
<b>IHC</b>	
<b>Reference</b>	Reference(s): de Haan G. Latexin is a newly discovered regulator of hematopoietic stem cells. Nature genetics 2007 Feb 39 (2): 141-2..PMID: 17262023->

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for **Research Use Only**