

Goat anti-SLC7A6 / y+LAT-2 Antibody

Item Number	dAP-3444
Target Molecule	Principle Name: SLC7A6 / y+LAT-2; Official Symbol: SLC7A6; All Names and Symbols: SLC7A6; solute carrier family 7 (amino acid transporter light chain, y+L system), member 6; LAT-2; LAT3; y+LAT-2; Y+L amino acid transporter 2; amino acid permease; cationic amino acid transporter, y+ system; solute carrier family 7 (cationic amino acid t; Accession Number (s): NP_003974.3; Human Gene ID(s): 9057; Non-Human GeneID(s):
Immunogen	PNTSQSQVEEDVSSP, is from N Terminus Reported variants represent identical protein: NP_003974.3, NP_001070253.1
Applications	Pep ELISA, WB Species Tested: Human
Purification	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Supplied As	Lyophilized powder of 50ug or 100ug IgG; Reconstitute IgG with 100ul or 200ul sterile DI Water and final product will be formulated as 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
Peptide ELISA	Peptide ELISA: antibody detection limit dilution 1 to 128000.
Western Blot	Western Blot: Approx 55-58kDa band observed in Human Placenta lysates (calculated MW of 56.8kDa according to NP_003974.3. Recommended concentration: 0.3-1µg/ml.
IHC	
Reference	Reference(s): Nguyen HT, Merlin D.Nguyen HT, Merlin D.Nguyen HT, Merlin D.Nguyen HT, Merlin D. Ho- meostatic and innate immune responses: role of the transmembrane glycoprotein CD98. Cellular and mo- lecular life sciences : CMLS 2012 Sep 69 (18): 3015-26..PMID: 22460579->

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the end users! This product is sold for **Research Use Only**