



## **Epithelial Neutrophil-Activating Protein 78 Mouse Recombinant (CXCL5)**

Item Number rAP-0133

Synonyms C-X-C motif chemokine 5, Small-inducible cytokine B5, Cytokine LIX, Cxcl5, Scyb5, LIX, GCP-2, Scyb6,

ENA-78, AMCF-II.

Description Epithelial Neutrophil-Activating Protein 78 Mouse Recombinant produced in E.Coli is a single, non-

glycosylated, polypeptide chain containing 93 amino acids and having a molecular mass of 9.8kDa. The

CXCL5 is purified by proprietary chromatographic techniques.

Uniprot Accesion Number P50228

Amino Acid Sequence APSSVIAATE LRCVCLTVTP KINPKLIANL EVIPAGPQCP TVEVIAKLKN QKEVCLDPEA PVIKKIIIQK

ILGSDKKKAK RNALAVERTA SVQ.

Source Escherichia Coli.

Physical Appearance

and Stability

Sterile Filtered White lyophilized (freeze-dried) powder. Lyophilized ENA-78 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution CXCL5 should be stored at 4°C between 2-7 days and for future use below -18°C.For long term storage it is recommended to

add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Formulation and Purity

Lyophilized from a 0.2µm filtered concentrated (1.0mg/ml) solution in 20mM sodium phosphate buffer, pH

7.4 & 150mM NaCl. Greater than 97.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-

PAGE.

**Application** 

Solubility It is recommended to reconstitute the lyophilized ENA-78 in sterile 18MΩ-cm H2O not less than 100μg/ml,

which can then be further diluted to other aqueous solutions.

Biological Activity Determined by its ability to chemoattract human peripheral blood neutrophils using a concentration range of

10-100ng/ml corresponding to a Specific Activity of 10,000-100,000IU/mg.

Shipping Format and Condition Lyophilized powder at room temperature.

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for Research Use Only