ANGIO-PROTEOMIE
HIGH QUALITY REAGENTS FOR LIFE SCIENCES

DATA SHEET

GRO-Gamma Human Recombinant (CXCL3)

Item Number	rAP-0157
Synonyms	Macrophage inflammatory protein 2-beta, MIP2-beta, CXCL3, Growth-regulated protein gamma, GRO-gamma, GRO-gamma(1-73), GRO3, GROg, MIP2B, SCYB3, MIP-2b, CINC-2b, MGSA gamma.
Description	GRO-Gamma Human Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 73 amino acids and having a molecular mass of 7902 Dalton. The CXCL3 is purified by proprietary chromatographic techniques.
Uniprot Accesion Number	P19876
Amino Acid Sequence	The sequence of the first five N-terminal amino acids was determined and was found to be Ala-Ser-Val-Val-Thr.
Source	Escherichia Coli.
Physical Appearance and Stability	Sterile Filtered White lyophilized (freeze-dried) powder. Lyophilized GRO-gamma although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution CXCL3 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Please prevent freeze-thaw cycles.
Formulation and Purity	The protein was lyophilized with no additives. Greater than 98.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.
Application	
Solubility	It is recommended to reconstitute the lyophilized CXCL3 in sterile 18MΩ-cm H2O not less than 100µg/ml, which can then be further diluted to other aqueous solutions.
Biological Activity	The Biological activity is calculated by its ability to chemoattract CXCR2 transfected 293 cells using 10-100ng/ml corresponding to a Specific Activity of 10,000-100,000IU/mg.
Shipping Format and Condition	Lyophilized powder at room temperature.

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for Research Use Only

Angio-Proteomie | 11 Park Drive, Suite 12 Boston, MA 02215, USA | Tel: 001-617-549-2665 | Fax: 001-480-247-4337 | Email: angioproteomie@gmail.com