

MIG Mouse Recombinant (CXCL9)

Item Number	rAP-0214
Synonyms	Small inducible cytokine B9, CXCL9, Gamma interferon-induced monokine, MIG, chemokine (C-X-C motif) ligand 9, CMK, Humig, SCYB9, crg-10, M119, monokine induced by gamma-interferon.
Description	MIG (monokine induced by gamma-interferon) Mouse Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 105 amino acids and having a molecular mass of 12208 Dalton. The MIG is purified by proprietary chromatographic techniques.
Uniprot Accesion Number	P18340
Amino Acid Sequence	The sequence of the first five N-terminal amino acids was determined and was found to be, Thr-Leu-Val-Ile-Arg.
Source	Escherichia Coli.
Physical Appearance and Stability	Sterile Filtered White lyophilized (freeze-dried) powder. Lyophilized MIG although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution CXCL9 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Please prevent freeze-thaw cycles.
Formulation and Purity	Filtered (0.2μm) and lyophilized from a concentrated (0.5mg/ml) solution in 20mM PB, pH7.4 and 100mM NaCl. Greater than 97.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.
Application	
Solubility	It is recommended to reconstitute the lyophilized MIG in sterile 18MΩ-cm H2O not less than 100μg/ml, which can then be further diluted to other aqueous solutions.
Biological Activity	The Activity is calculated by the ability to chemoattract Human lymphocytes using a concentration of 0.1-1 ng/ml.
Shipping Format and Condition	Lyophilized powder at room temperature.

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for **Research Use Only**