

## Interleukin 17F Human Recombinant

<b>Item Number</b>	rAP-0617
<b>Synonyms</b>	Cytokine ML-1, IL-17F, Interleukin-17F precursor, IL17F, ML1, ML-1.
<b>Description</b>	Interleukin-17F Human Recombinant produced in E.Coli is a homodimeric, cysteine linked, non-glycosylated polypeptide chain containing 2 x 134 amino acids and having a total molecular mass of 30.1 kDa. The IL-17F Human is purified by proprietary chromatographic techniques.
<b>Uniprot Accession Number</b>	Q96PD4
<b>Amino Acid Sequence</b>	MRKIPKVGHT FFQKPESCPP VPGGSMKLDI GIINENQRVS MSRNIESRST SPWNYTVTWD PNRYPSEVVQ AQCRNLGCIN AQGKEDISMN SVPIQQETLV VRRKHQGCSV SFQLEKVLVT VGCTCVTPVI HHVQ.
<b>Source</b>	Escherichia Coli.
<b>Physical Appearance and Stability</b>	Sterile Filtered White lyophilized (freeze-dried) powder. Lyophilized Interleukin Human IL17F although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution Human IL-17F should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.
<b>Formulation and Purity</b>	Lyophilized from a concentrated (1mg/ml) solution containing 10mM sodium citrate buffer pH-3. Greater than 97.0% as determined by (a) Analysis by RP-HPLC. (b) Analysis by SDS-PAGE.
<b>Application</b>	
<b>Solubility</b>	It is recommended to reconstitute the lyophilized Interleukin Human IL17F in sterile water not more than 1mg/ml, which can then be further diluted to other aqueous solutions.
<b>Biological Activity</b>	
<b>Shipping Format and Condition</b>	Lyophilized powder at room temperature.

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for **Research Use Only**