

DNA Damage Inducible Transcript 3 Human Recombinant

Item Number	rAP-3197
Synonyms	DNA damage-inducible transcript 3, DDIT-3, Growth arrest and DNA-damage-inducible protein GADD153, C/EBP-homologous protein, CHOP, DDIT3, GADD153, CEBPZ, CHOP10, MGC4154.
Description	DDIT3 Human Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 189 amino acids and having a molecular mass of 21 kDa. The DDIT3 protein is fused to a 20 amino acids His tag at N-terminus. The DDIT3 is purified by proprietary chromatographic techniques.
Uniprot Accession Number	P35638
Amino Acid Sequence	MGSSHHHHHH SSGLVPRGSH MAAESLPFSF GTLSSWELEA WYEDLQEVLS SDENGGTYVS PPGN- EEEEESK IFTTLDPASL AWLTEEEPEP AEVTSTSQSP HSPDSSQSSL AQEEEEEDQG RTRKRKQSGH SPARAGKQRM KEKEQENERK VAQLAEENER LKQEIERTLR EVEATRRALI DRMVNLHQA.
Source	Escherichia Coli.
Physical Appearance and Stability	Sterile Filtered colorless solution. Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.
Formulation and Purity	The DDIT3 protein solution (1mg/ml) contains 20mM Tris-HCl pH-8 and 20% glycerol. Greater than 90.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.
Application	
Solubility	
Biological Activity	
Shipping Format and Condition	Lyophilized powder at room temperature.

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for **Research Use Only**