

## Mouse Monoclonal Antibody to CD10

Catalogue Number	sAP-0209
Target Molecule	<b>Name:</b> CD10 <b>Aliases:</b> NEP; CALLA <b>MW:</b> N/A
Description	CD10(MME): membrane metallo-endopeptidase. This gene encodes a common acute lymphocytic leukemia antigen that is an important cell surface marker in the diagnosis of human acute lymphocytic leukemia (ALL). This protein is present on leukemic cells of pre-B phenotype, which represent 85% of cases of ALL. This protein is not restricted to leukemic cells, however, and is found on a variety of normal tissues. It is a glycoprotein that is particularly abundant in kidney, where it is present on the brush border of proximal tubules and on glomerular epithelium. The protein is a neutral endopeptidase that cleaves peptides at the amino side of hydrophobic residues and inactivates several peptide hormones including glucagon, enkephalins, substance P, neurotensin, oxytocin, and bradykinin. This gene, which encodes a 100-kD type II trans-
Immunogen	Purified recombinant fragment of CD10 expressed in E. Coli. ;
Recitative Species	Human
Clone	MM3G9D10;
Size and Concentration	100µg/1mg/ml
Supplied as	Lyophilized Powder from 100µl of Ascitic fluid containing 0.03% sodium azide. ;
Reconstitution/Storages	Reconstituted with 100µl sterile DI H2O, at stored at 4°C or -20°C for short or long term storage
Applications	ELISA: 1 to 10000; WB: 1 to 500 - 1 to 2000; IHC: 1 to 200 - 1 to 1000
Shipping	Regular FEDEX overnight shipment (ambient temperature)
Reference	1. Journal of Gastroenterology, 1996 Feb.31(1):12-7. ; 2. British Journal of Haematology, 1995,89(3):623-6.

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for **Research Use Only**