

## Mouse Monoclonal Antibody to HK1

<b>Catalogue Number</b>	sAP-0442
<b>Target Molecule</b>	<b>Name:</b> HK1 <b>Aliases:</b> HK1; HXK1; HK1-ta; HK1-tb; HK1-tc; HK1 <b>MW:</b> 102kDa <b>Entrez Gene ID:</b> 3098
<b>Description</b>	The hexokinases utilize Mg-ATP as a phosphoryl donor to catalyze the first step of intracellular glucose metabolism, the conversion of glucose to glucose- 6-phosphate. Four hexokinase isoenzymes have been identified, including hexokinase I (HXK I), hexokinase II (HXK II), hexokinase III (HXK III) and hexokinase IV (HXK IV, also designated glucokinase or GCK). Hexokinases I-III each contain an N-terminal cluster of hydrophobic amino acids. Glucokinase lacks the N-terminal hydrophobic cluster. The hydrophobic cluster is thought to be necessary for membrane binding. This is substantiated by the finding that glucokinase has lower affinity for glucose than do the other hexokinases. HK I has been shown to be expressed in brain, kidney and heart tissues as well as in hepatoma cell lines.
<b>Immunogen</b>	Purified recombinant fragment of human HK1 expressed in E. Coli.
<b>Recitative Species</b>	Human; Mouse; Rat;
<b>Clone</b>	MM3A10;
<b>Size and Concentration</b>	100 $\mu$ g/1mg/ml
<b>Supplied as</b>	Lyophilized Powder from 100 $\mu$ l of Ascitic fluid containing 0.03% sodium azide.
<b>Reconstitution/Storages</b>	Reconstituted with 100 $\mu$ l sterile DI H <sub>2</sub> O, at stored at 4°C or -20°C for short or long term storage
<b>Applications</b>	ELISA: 1 to 10000; WB: 1 to 500 - 1 to 2000; IHC: 1 to 200 - 1 to 1000; ICC: 1 to 200 - 1 to 1000; FCM: 1 to 200 - 1 to 400
<b>Shipping</b>	Regular FEDEX overnight shipment (ambient temperature)
<b>Reference</b>	1. Cell. 2005 Sep 23;122(6):957-68.; 2. Am J Hum Genet. 2006 Jan;78(1):78-88.

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for **Research Use Only**