

## Mouse Monoclonal Antibody to Oct4

<b>Catalogue Number</b>	sAP-0471
<b>Target Molecule</b>	<b>Name: Oct4</b> <b>Aliases:</b> OCT3; OCT4; OTF3; OTF4; OTF-3; Oct-3; Oct-4; MGC22487; POU5F1 <b>MW: 45kDa</b> <b>Entrez Gene ID: 5460</b>
<b>Description</b>	This gene encodes a transcription factor containing a POU homeodomain. This transcription factor plays a role in embryonic development, especially during early embryogenesis, and it is necessary for embryonic stem cell pluripotency. A translocation of this gene with the Ewing's sarcoma gene, t(6;22)(p21;q12), has been linked to tumor formation. Alternative splicing, as well as usage of alternative translation initiation codons, results in multiple isoforms, one of which initiates at a non-AUG (CUG) start codon. Related pseudogenes have been identified on chromosomes 1, 3, 8, 10, and 12. (provided by RefSeq). Tissue specificity: Expressed in developing brain. Highest levels found in specific cell layers of the cortex, the olfactory bulb, the hippocampus and the cerebellum. Low levels of expression in adult tissues.
<b>Immunogen</b>	Synthesized peptide derived from internal of human Oct4.
<b>Recitative Species</b>	Human
<b>Clone</b>	MM7E7;
<b>Size and Concentration</b>	100µg/1mg/ml
<b>Supplied as</b>	Lyophilized Powder from 100µl of Ascitic fluid containing 0.03% sodium azide.
<b>Reconstitution/Storages</b>	Reconstituted with 100µl sterile DI H <sub>2</sub> O, at stored at 4°C or -20°C for short or long term storage
<b>Applications</b>	ELISA: 1 to 10000; WB: 1 to 500 - 1 to 2000; ICC: 1 to 200 - 1 to 1000; FCM: 1 to 200 - 1 to 400
<b>Shipping</b>	Regular FEDEX overnight shipment (ambient temperature)
<b>Reference</b>	1. Stem Cells. 2010 May;28(5):885-93. ; 2. Mol Med. 2010 Jul-Aug;16(7-8):247-53. ; 3. Med Sci (Paris). 2010 Apr;26(4):411-6.

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for **Research Use Only**