

Mouse Monoclonal Antibody to NFKB1

Catalogue Number	sAP-0574
Target Molecule	<p>Name: NFKB1</p> <p>Aliases: p50; KBF1; p105; EBP-1; MGC54151; NFKB-p50; NFkappaB; NF-kappaB; NFKB-p105; NF-kappa-B</p> <p>MW: 50kDa/105kDa</p>
Description	<p>This gene encodes a 105 kD protein which can undergo cotranslational processing by the 26S proteasome to produce a 50 kD protein. The 105 kD protein is a Rel protein-specific transcription inhibitor and the 50 kD protein is a DNA binding subunit of the NF-kappa-B (NFKB) protein complex. NFKB is a transcription regulator that is activated by various intra- and extra-cellular stimuli such as cytokines, oxidant-free radicals, ultraviolet irradiation, and bacterial or viral products. Activated NFKB translocates into the nucleus and stimulates the expression of genes involved in a wide variety of biological functions. Inappropriate activation of NFKB has been associated with a number of inflammatory diseases while persistent inhibition of NFKB leads to inappropriate immune cell development or delayed cell growth. Two transcript variants en-</p>
Immunogen	Purified recombinant fragment of human NFKB1 expressed in E. Coli. ;
Recitative Species	Human
Clone	MM5D10;
Size and Concentration	100µg/1mg/ml
Supplied as	Lyophilized Powder from 100µl of Ascitic fluid containing 0.03% sodium azide.
Reconstitution/Storages	Reconstituted with 100µl sterile DI H ₂ O, at stored at 4°C or -20°C for short or long term storage
Applications	ELISA: 1 to 10000; WB: 1 to 500 - 1 to 2000; IHC: 1 to 200 - 1 to 1000; FCM: 1 to 200 - 1 to 400
Shipping	Regular FEDEX overnight shipment (ambient temperature)
Reference	1. Cytokine. 2010 Feb;49(2):215-20. ; 2. Chemotherapy. 2009;55(5):381-5.

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for **Research Use Only**