

Mouse Monoclonal Antibody to CD22

Catalogue Number	sAP-0844
Target Molecule	Name: CD22 Aliases: SIGLEC2; SIGLEC-2 MW: 95.3kDa Entrez Gene ID: 933
Description	CD22 may be involved in the localization of B-cells in lymphoid tissues. Binds sialylated glycoproteins; one of which is CD45. Preferentially binds to alpha-2,6-linked sialic acid. The sialic acid recognition site can be masked by cis interactions with sialic acids on the same cell surface. Upon ligand induced tyrosine phosphorylation in the immune response seems to be involved in regulation of B-cell antigen receptor signaling. Plays a role in positive regulation through interaction with Src family tyrosine kinases and may also act as an inhibitory receptor by recruiting cytoplasmic phosphatases via their SH2 domains that block signal transduction through dephosphorylation of signaling molecules ;
Immunogen	Purified recombinant fragment of human CD22 (AA: 621-725) expressed in E. Coli.
Reactive Species	Human;
Clone	MM2H1C4;
Size and Concentration	100µg/1mg/ml
Supplied as	Lyophilized Powder from 100µl of Purified antibody in PBS with 0.05% sodium azide
Reconstitution/Storages	Reconstituted with 100µl sterile DI H ₂ O, at stored at 4°C or -20°C for short or long term storage
Applications	ELISA: 1 to 10000; WB: 1 to 500 - 1 to 2000; IHC: 1 to 200 - 1 to 1000; ICC: 1 to 200 - 1 to 1000; FCM: 1 to 200 - 1 to 400
Shipping	Regular FEDEX overnight shipment (ambient temperature)
Reference	1. Cancer Res. 2012 Nov 1;72(21):5556-65. ; 2. J Innate Immun. 2011;3(4):411-9. ;

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for **Research Use Only**