

## Mouse Monoclonal Antibody to NR1I2

<b>Catalogue Number</b>	sAP-1448
<b>Target Molecule</b>	<p><b>Name:</b> NR1I2</p> <p><b>Aliases:</b> APC11; Apc11p; HSPC214</p> <p><b>MW:</b> 9.8kDa</p> <p><b>Entrez Gene ID:</b> 51529</p>
<b>Description</b>	<p>This gene product belongs to the nuclear receptor superfamily, members of which are transcription factors characterized by a ligand-binding domain and a DNA-binding domain. The encoded protein is a transcriptional regulator of the cytochrome P450 gene CYP3A4, binding to the response element of the CYP3A4 promoter as a heterodimer with the 9-cis retinoic acid receptor RXR. It is activated by a range of compounds that induce CYP3A4, including dexamethasone and rifampicin. Several alternatively spliced transcripts encoding different isoforms, some of which use non-AUG (CUG) translation initiation codon, have been described for this gene. Additional transcript variants exist, however, they have not been fully characterized.;</p>
<b>Immunogen</b>	Purified recombinant fragment of human NR1I2 (AA:1-142) expressed in E. Coli.
<b>Reactive Species</b>	Human;
<b>Clone</b>	MM6H11D8;
<b>Size and Concentration</b>	100µg/1mg/ml
<b>Supplied as</b>	Lyophilized Powder from 100µl of Purified antibody in PBS with 0.05% sodium azide
<b>Reconstitution/Storages</b>	Reconstituted with 100µl sterile DI H <sub>2</sub> O, at stored at 4°C or -20°C for short or long term storage
<b>Applications</b>	ELISA: 1 to 10000; WB: 1 to 500 - 1 to 2000; IHC: ; ICC: ; FCM: 1 to 200 - 1 to 400
<b>Shipping</b>	Regular FEDEX overnight shipment (ambient temperature)
<b>Reference</b>	1.Int J Mol Sci. 2014 Sep 29;15(10):17457-68. ; 2.J Periodontal Res. 2012 Apr;47(2):174-9. ;

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for **Research Use Only**