

Mouse Monoclonal Antibody to IGHM

Catalogue Number	sAP-1567
Target Molecule	Name: IGHM Aliases: MU; VH; AGM1 MW: 49.3kDa Entrez Gene ID: 3507
Description	Immunoglobulins (Ig) are the antigen recognition molecules of B cells. An Ig molecule is made up of 2 identical heavy chains and 2 identical light chains (see MIM 147200) joined by disulfide bonds so that each heavy chain is linked to a light chain and the 2 heavy chains are linked together. Each Ig heavy chain has an N-terminal variable (V) region containing the antigen-binding site and a C-terminal constant (C) region, encoded by an individual C region gene, that determines the isotype of the antibody and provides effector or signaling functions. The heavy chain V region is encoded by 1 each of 3 types of genes: V genes (see MIM 147070), joining (J) genes (see MIM 147010), and diversity (D) genes (see MIM 146910). The C region genes are clustered downstream of the V region genes within the heavy chain locus on chromosome 14.
Immunogen	Purified recombinant fragment of human IGHM (AA: 310-452) expressed in E. Coli.
Recitative Species	Human;
Clone	MM9A8A4
Size and Concentration	100µg/1mg/ml
Supplied as	Lyophilized Powder from 100µl of Purified antibody in PBS with 0.05% sodium azide
Reconstitution/Storages	Reconstituted with 100µl sterile DI H2O, at stored at 4°C or -20°C for short or long term storage
Applications	ELISA: 1 to 10000; WB: 1 to 500 - 1 to 2000; ICC: N to A; FCM: 1 to 200 - 1 to 400; IHC: 1 to 200 - 1 to 1000
Shipping	Regular FEDEX overnight shipment (ambient temperature)
Reference	1. Cell Mol Immunol. 2014 Jan;11(1):94-104. 2. Mol Immunol. 1993 Jan;30(1):111-2.

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the end users! This product is sold for **Research Use Only**