

Mouse Monoclonal Antibody to BTRC

Catalogue Number	sAP-1613
Target Molecule	Name: BTRC Aliases: FWD1; FBW1A; FBXW1; bTrCP; FBXW1A; bTrCP1; betaTrCP; BETA-TRCP MW: 68.9kDa
Description	Entrez Gene ID: 8945 This gene encodes a member of the F-box protein family which is characterized by an approximately 40 amino acid motif, the F-box. The F-box proteins constitute one of the four subunits of ubiquitin protein ligase complex called SCFs (SKP1-cullin-F-box), which function in phosphorylation-dependent ubiquitination. The F-box proteins are divided into 3 classes: Fbws containing WD-40 domains, Fbls containing leucine-rich repeats, and Fbxs containing either different protein-protein interaction modules or no recognizable motifs. The protein encoded by this gene belongs to the Fbws class; in addition to an F-box, this protein contains multiple WD-40 repeats. The encoded protein mediates degradation of CD4 via its interaction with HIV-1 Vpu. It has also been shown to ubiquitinate phosphorylated NFKBIA (nuclear factor of kappa light
Immunogen	Purified recombinant fragment of human BTRC (AA: 24-151) expressed in E. Coli.
Recitative Species	Human;
Clone	MM3D5E6
Size and Concentration	100µg/1mg/ml
Supplied as	Lyophilized Powder from 100µl of Purified antibody in PBS with 0.05% sodium azide
Reconstitution/Storages	Reconstituted with 100µl sterile DI H2O, at stored at 4°C or -20°C for short or long term storage
Applications	ELISA: 1 to 10000; WB: 1 to 500 - 1 to 2000; ICC: N to A; FCM: N to A; IHC: 1 to 200 - 1 to 1000
Shipping	Regular FEDEX overnight shipment (ambient temperature)
Reference	1.Biochem Biophys Res Commun. 2013 Nov 29;441(4):831-7.2.PLoS One. 2011;6(11):e27464.

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for **Research Use Only**