

Mouse Monoclonal Antibody to CD197

Catalogue Number	sAP-1677
Target Molecule	Name: CD197 Aliases: BLR2; EBI1; CCR-7; CCR7; CDw197; CMKBR7; CC-CKR-7 MW: 42.9kDa Entrez Gene ID: 1236
Description	The protein encoded by this gene is a member of the G protein-coupled receptor family. This receptor was identified as a gene induced by the Epstein-Barr virus (EBV), and is thought to be a mediator of EBV effects on B lymphocytes. This receptor is expressed in various lymphoid tissues and activates B and T lymphocytes. It has been shown to control the migration of memory T cells to inflamed tissues, as well as stimulate dendritic cell maturation. The chemokine (C-C motif) ligand 19 (CCL19/ECL) has been reported to be a specific ligand of this receptor. Signals mediated by this receptor regulate T cell homeostasis in lymph nodes, and may also function in the activation and polarization of T cells, and in chronic inflammation pathogenesis. Alternative splicing of this gene results in multiple transcript variants.
Immunogen	Purified recombinant fragment of human CD197 (AA: extra mix) expressed in E. Coli.
Recitative Species	Human;
Clone	MM7A5A9
Size and Concentration	100µg/1mg/ml
Supplied as	Lyophilized Powder from 100µl of Purified antibody in PBS with 0.05% sodium azide
Reconstitution/Storages	Reconstituted with 100µl sterile DI H2O, at stored at 4°C or -20°C for short or long term storage
Applications	ELISA: 1 to 10000; WB: 1 to 500 - 1 to 2000; ICC: N to A; FCM: 1 to 200 - 1 to 400; IHC: N to A
Shipping	Regular FEDEX overnight shipment (ambient temperature)
Reference	1.Gastric Cancer. 2017 Mar;20(2):235-245.2.J Surg Oncol. 2015 Jul;112(1):86-92.

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for **Research Use Only**