

## Mouse Monoclonal Antibody to CD209

<b>Catalogue Number</b>	sAP-1687
<b>Target Molecule</b>	<b>Name:</b> CD209 <b>Aliases:</b> CDSIGN; CLEC4L; DC-SIGN; DC-SIGN1 <b>MW:</b> 45.8kDa <b>Entrez Gene ID:</b> 30835
<b>Description</b>	This gene encodes a transmembrane receptor and is often referred to as DC-SIGN because of its expression on the surface of dendritic cells and macrophages. The encoded protein is involved in the innate immune system and recognizes numerous evolutionarily divergent pathogens ranging from parasites to viruses with a large impact on public health. The protein is organized into three distinct domains: an N-terminal transmembrane domain, a tandem-repeat neck domain and C-type lectin carbohydrate recognition domain. The extracellular region consisting of the C-type lectin and neck domains has a dual function as a pathogen recognition receptor and a cell adhesion receptor by binding carbohydrate ligands on the surface of microbes and endogenous cells. The neck region is important for homo-oligomerization which allows the receptor to bind to two different ligands.
<b>Immunogen</b>	Purified recombinant fragment of human CD209 (AA: extra 270-404) expressed in E. Coli.
<b>Recitative Species</b>	Human;
<b>Clone</b>	MM5C2A6
<b>Size and Concentration</b>	100µg/1mg/ml
<b>Supplied as</b>	Lyophilized Powder from 100µl of Purified antibody in PBS with 0.05% sodium azide
<b>Reconstitution/Storages</b>	Reconstituted with 100µl sterile DI H2O, at stored at 4°C or -20°C for short or long term storage
<b>Applications</b>	ELISA: 1 to 10000; WB: 1 to 500 - 1 to 2000; ICC: N to A; FCM: 1 to 200 - 1 to 400; IHC: N to A
<b>Shipping</b>	Regular FEDEX overnight shipment (ambient temperature)
<b>Reference</b>	1.Blood. 2015 Oct 15;126(16):1911-20. 2.PLoS One. 2014 Aug 22;9(8):e105236.

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the end users! This product is sold for **Research Use Only**